

CLAIMS

We claim:

1. An isolated polypeptide of at least 15 amino acid residues comprising an epitope-bearing portion of a protein of SEQ ID NO:2 or SEQ ID NO:43.

2. The isolated polypeptide according to claim 1 wherein said polypeptide is selected from the group consisting of:

residues 15-163 of SEQ ID NO:2;
 residues 46-163 of SEQ ID NO:2;
 residues 15-170 of SEQ ID NO:2;
 residues 46-170 of SEQ ID NO:2;
 residues 15-234 of SEQ ID NO:2;
 residues 46-234 of SEQ ID NO:2;
 residues 15-229 amide of SEQ ID NO:2;
 residues 15-230 of SEQ ID NO:2;
 residues 15-345 of SEQ ID NO:2;
 residues 46-345 of SEQ ID NO:2;
 residues 164-345 of SEQ ID NO:2;
 residues 235-345 of SEQ ID NO:2; and
 residues 226-345 of SEQ ID NO:2.

3. An isolated polypeptide comprising a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein:

R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2;

R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2;

R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2; a glycine residue

007220: 007220

at a position corresponding to residue 282 of SEQ ID NO:2; and a sequence motif CX{18,33}CXGXCX{6,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:

at least one of x and z is 1; and
if x and z are each 1, then y is 1.

4. The isolated polypeptide of claim 3 wherein x=1.

5. The isolated polypeptide of claim 4 wherein R1 comprises residues 46-163 of SEQ ID NO:2.

6. The isolated polypeptide of claim 4 wherein R1 is at least 90% identical to residues 18-163 of SEQ ID NO:2.

7. The isolated polypeptide of claim 4 wherein y=1.

8. The isolated polypeptide of claim 7 wherein z=1.

9. The isolated polypeptide of claim 3 wherein said polypeptide comprises residues 46-229 of SEQ ID NO:2, residues 164-345 of SEQ ID NO:2, or residues 46-345 of SEQ ID NO:2.

10. The isolated polypeptide of claim 3 wherein z=1.

11. The isolated polypeptide of claim 10 wherein R3 comprises residues 235-345 of SEQ ID NO:2.

12. The isolated polypeptide of claim 10 wherein y=1.

007220 35274560

13. The isolated polypeptide of claim 12 wherein $x=1$.

14. The isolated polypeptide of claim 10, further comprising cysteine residues at positions corresponding to residues 286, 287, 291, and 294 of SEQ ID NO:2.

15. The isolated polypeptide of claim 3, further comprising an affinity tag.

16. The isolated polypeptide of claim 3, further comprising an immunoglobulin constant domain.

17. An isolated protein comprising a first polypeptide operably linked to a second polypeptide, wherein said first polypeptide comprises a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein:

R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[ER][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2;

R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2;

R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2; and a sequence motif CX{25,33}CXGXCX{10,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:

at least one of x and z is 1; and

if x and z are each 1, then y is 1,

and wherein said protein modulates cell proliferation, differentiation, metabolism, or migration.

cysteine residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2; and a sequence motif CX{25,33}CXGXCX{10,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and

at least one of x and z is 1; and
if x and z are each 1, then y is 1.

surface PDGF alpha receptor to the protein of claim 17, whereby the protein binds to and activates the receptor.

40. A method of inhibiting a PDGF alpha receptor-mediated cellular process, comprising exposing a cell comprising a cell-surface PDGF alpha receptor to a compound that inhibits binding of the protein of claim 17 to the receptor.

41. A method of inhibiting zveg3 activity in a mammal comprising administering to the mammal an effective amount of a zveg3 antagonist.

42. The method of claim 41 wherein the antagonist is an antibody, a receptor, a ligand-binding receptor fragment, or a receptor IgG-Fc fusion protein.

43. An isolated, antisense polynucleotide that is the complement of the isolated polynucleotide of claim 25.

44. The antisense polynucleotide of claim 43 further comprising operably linked transcription promoter and terminator sequences.

45. A method of inhibiting zveg3 production in a cell comprising administering to the cell the antisense polynucleotide of claim 44.

00760 2327500